



CRYSTAL CLEAR
SKINCARE

Intense Pulse Light Therapy

Hair Removal Price List

Area	Single Session Pay as you go	Course of 10 Total amount paid upfront
Consultation and Patch Test	FREE	-
Upper Lip	£35.00	£280
Chin	£35.00	£280
Side Burns / Full Face	£55.00 / £65.00	£440 / £520
Underarms	£55.00	£440
Bikini Line	£70.00	£560
Brazilian	£100.00	£800
Hollywood	£150.00	£1200
Half Leg	£180.00	£1440
³/₄ Leg	£210.00	£1680
Full Leg	£250.00	£2000
Full Back	£200.00	£1600
Full Chest	£150.00	£1200

Treatments are taken in four to eight week intervals, and a course of eight to ten treatments is recommended.

A Patch Test will be undertaken prior to treatment to ascertain suitability for treatment.

Please bring any medication you are taking with you to this appointment as some medications are contraindicated to the treatment.

See Contra-Indications/ Medicines list

The following points need to be considered:

- **Treatment is not possible if you have a tan, (natural, fake or from a sunbed).**
- **Treatment is not possible if you are pregnant.**
- **Treatment is not possible if you have suffered from any type of skin cancer in the last 10 years.**
- **You will be advised to refrain from any strenuous exercise or any heat treatments for 48 hours following the patch test.**
- **For hair removal we will need to have a small area shaved the day before consultation, but do not shave the entire area as we need to assess hair type.**
- **One or two days before a full treatment the whole area will need to be carefully shaved, this is very important as any hair left unshaved can cause burning on the skin. We do not have the facility to shave hair so this must be done before attending your appointment.**
- **Following a Patch Test your course of treatment must be undertaken within at least 6 months or another patch test will be required. If any change in medical condition or medication occurs after the patch test has been performed we may need to re-test before proceeding with treatment.**
- **In order to qualify for the reduced prices of a course, the full course price must be paid before commencing treatment.**

Contra-indications

The contra-indications for the IPL procedure are minimal, however the following exclusion from treatment criteria is recommended:

- Tanned skin (active tan) including Fake tan
- Pregnancy (do not treat until first period and end of breastfeeding)
- Hepatitis
- Heart Conditions (pacemaker)
- Porphyria (allergies to sunlight)
- Auto immune diseases (Lupus, aids, HIV etc)
- A history of keloid scarring;
- Liver or kidney disease
- Herpes (shingles/cold sores)
- Use of medication that may induce photosensitivity to the skin (e.g hydrocortisone) or perfume products, deodorants, sun block, essential oils;
- Any inflammatory skin condition at the treatment site;
- A history of cancer including cancer medication, radiation and chemotherapy
- Abnormal swelling or oedema
- Botox, collagen or restylane within last 6 months in area to be treated
- The client has a history of poor wound healing including Type I Diabetes.
- Vitiligo (loss of pigmentation within the skin)
- Treatment over certain skin areas such as tattoos, moles, semi permanent make-up, lip vermilion or mucous membranes.
- Anti-Coagulants (blood thinning medication)
- Accutane (given by doctors for acne)
- Isotretinoin, Tretinoin, Retin A in previous 3-6 months as can lead to thinning of the skin
- Anyone under 18 years of age if establishment is not registered to treat children or is parental permission is with held.

IPL After Care Advice

- Don't expose skin to UV (Sun, Sunbeds and fake tan) for at least 2 weeks
- Don't shave for 48-72 hrs after treatment
- Don't depilate(wax, pluck, threading, tweeze) between treatments
- Don't use bleaching creams or perfumed products for 24 - 48hrs
- Don't pick or scratch the area
- Avoid rough handling of the area
- Leave any skin response alone these are temporary and will subside

- Metal Pins or plates in the area to be treated
- Epilepsy
- St Johns Wart (herbal medication) should not be used for at least 3 months prior to the treatment as can lead to photosensitivity
- Waxing, plucking, sugaring or threading in the area to be treated 4- 6weeks prior as the hair needs to be in the follicle for the treatment to work.
- **Black, Afro Caribbean skin, very dark skin NOT TREATABLE WITH IPL**

Common Drugs That Can Cause Photosensitive Reactions

- Antibiotics
- Dioxycycline (Vibramycin, Vibratabs)
- Demeclocycline (Declomycin) – High incidence.
- Tetracycline (Achromycin and others)
- Nalidixic Acid (NegGram)
- Lomefloxacin (Maxaquin) – Especially noted for severe reactions when they occur.

Blood Pressure and Heart Medications

- Hydrochlorothiazide (Hydrodiuril, Oretic and other names) – Many drugs contain HCTZ as one of the ingredients.
- Chlorothiazide (Diuril and other names)
- Furosemide (Lasix)
- Amiodarone (Cordarone) - High incidence of drug-induced photoreactions.

Other Drugs

- Chlorpromazine (Thorazine)
- Methoxypsoralin
- PABA and/or PABA esters – Used in sunscreens, can cause hives and rashes.

- **Avoid very hot baths, steam treatments saunas for 1 week**
- **Avoid swimming for 1 week**
- **Avoid exfoliating, peels and Microdermabrasion for 1 week**
- **Avoid rough sports for 24-48hrs**
- **Avoid wearing tight clothing**
- **Keep the area clean and dry**
- **Keep hydrated**
- **Use a sun block (Crystal Clear Protect and Repair SPF40 recommended**

If an adverse reaction was to occur apply cold compressor ice pack to the area within 24 hours. Contact the salon ASAP leanne_beauty@hotmail.co.uk 01279 871004